

Maximum Probability and Maximum Entropy methods: Bayesian interpretation

M. Grendar, Jr. and M. Grendar

*Institute of Mathematics and Computer Science of Mathematical Institute of Slovak Academy of Sciences (SAS) and of Matej Bel University, Severná ulica 5, 974 00 Banská Bystrica, Slovakia & Institute of Measurement Science of SAS, Dúbravská cesta 9, 841 04 Bratislava, Slovakia.
umergren@savba.sk*

Abstract. (Jaynes') Method of (Shannon-Kullback's) Relative Entropy Maximization (REM or MaxEnt) can be - at least in the discrete case - according to the Maximum Probability Theorem (MPT) viewed as an asymptotic instance of the Maximum Probability method (MaxProb). A simple bayesian interpretation of MaxProb is given here. MPT carries the interpretation over into REM.

INTRODUCTION

Relationship of the Method of (Shannon-Kullback's) Relative Entropy Maximization (REM or MaxEnt) and Bayesian Method is notoriously peculiar. The two methods of induction are viewed as unrelated at all, or opposed, or identical in some circumstances, or one as a special case of the other one (see [8]).

As it was noted, a finding that REM can be viewed as an asymptotic instance of Maximum Probability method (MaxProb, cf. [3]) implies that MaxProb/REM/MaxEnt cannot be in conflict with Bayes' Theorem (cf. [4]).

A beautiful, simple (yet in some extent overlooked) bayesian interpretation of REM which operates on the level of samples and employs Conditioned Weak Law of Large Numbers (CWLLN) was suggested and elaborated at [2]. Csizsár's original argument together with the Maximum Probability Theorem (MPT, see [3], Thm 1), inspired a bayesian interpretation of MaxProb and REM methods, which we intend to present here.

TERMINOLOGY AND NOTATION

Let $\mathcal{X} \triangleq \{x_1, x_2, \dots, x_m\}$ be a discrete finite set called support, with m elements and let $\{X_l, l = 1, 2, \dots, n\}$ be a sequence of size n of identically and independently drawn random variables taking values in $\mathcal{X}$.

A type $\mathbf{v}_n \triangleq [n_1, n_2, \dots, n_m]/n$ is an empirical probability mass function which can be based on se-

quence $\{X_l, l = 1, 2, \dots, n\}$. Thus, n_i denotes number of occurrences of i -th element of $\mathcal{X}$ in the sequence.

Let $\mathcal{P}(\mathcal{X})$ be a set of all probability mass functions (pmf's) on $\mathcal{X}$. Let $\Pi_n \subseteq \mathcal{P}(\mathcal{X})$ be a set of all types $\mathbf{v}_n$, and let $\mathcal{H}_n \subseteq \Pi_n$.

Let the supposed source of the sequences (and hence also of types) be $\mathbf{q} \in \mathcal{P}(\mathcal{X})$.

Let $\pi(\mathbf{v}_n)$ denote the probability that $\mathbf{q}$ will generate type $\mathbf{v}_n$, ie. $\pi(\mathbf{v}_n) \triangleq \frac{n!}{n_1! n_2! \dots n_m!} \prod_{i=1}^m q_i^{n_i}$.

BAYESIAN INTERPRETATION OF MAXIMUM PROBABILITY METHOD

Bayesian recipe prescribes to update prior distribution (information) by an evidence via Bayes' Theorem (BT) to get a posterior distribution. Usually bayesians use BT to update prior distribution of a parameter by evidence which has form of random sample and obtain posterior distribution of the parameter, given the sample. Then it is customary to select the value of parameter at which the posterior distribution attains its maximum (i.e. mode) and perform further inference.

The bayesian recipe and [2] will be followed here on a different level. A prior distribution of types will be updated via BT by data of special form. Then the maximum aposteriori type will be searched out.

The bayesian updating will be carried out in four steps:

Step 1: Select a probability mass function $\mathbf{q}$ which

could be the best guess of source of types $\mathbf{v}_n$. It will specify a prior probability $P(\mathbf{v}_n)$ of type by the following simple scheme: $P(\mathbf{v}_n) \equiv \pi(\mathbf{v}_n)$. Thus, $\pi(\mathbf{v}_n)$ is the apriori distribution of types, which is going to be updated once an evidence (data) will become available.

Step 2: The data arrive in rather special form: they specify a set $\mathcal{H}_n$ of types $\mathbf{v}_n$ (which were observed, or 'feasible' in some general way). In other words, the evidence is that types which do not belong to $\mathcal{H}_n$ cannot be observed, or are 'not feasible'.

Step 3: Use Bayes' Theorem to update the prior probability of type $\pi(\text{type} = \mathbf{v}_n)$ by the evidence "type $\in \mathcal{H}_n$ " to obtain the posterior probability $P(\text{type} = \mathbf{v}_n | \text{type} \in \mathcal{H}_n)$ that type is equal to $\mathbf{v}_n$ given that it conforms with the evidence (i.e. belongs to $\mathcal{H}_n$).

$$P(\text{type} = \mathbf{v}_n | \text{type} \in \mathcal{H}_n) = \frac{P(\text{type} \in \mathcal{H}_n | \text{type} = \mathbf{v}_n) \pi(\text{type} = \mathbf{v}_n)}{P(\text{type} \in \mathcal{H}_n)}$$

Note that $P(\text{type} \in \mathcal{H}_n | \text{type} = \mathbf{v}_n)$ is 0 if $\mathbf{v}_n \notin \mathcal{H}_n$ and 1 otherwise. Thus, for $\mathbf{v}_n \in \mathcal{H}_n$ the aposteriori probability is

$$P(\text{type} = \mathbf{v}_n | \text{type} \in \mathcal{H}_n) = \frac{\pi(\text{type} = \mathbf{v}_n)}{P(\text{type} \in \mathcal{H}_n)}$$

Obviously, $P(\text{type} \in \mathcal{H}_n) = \sum_{\mathbf{v}_n \in \mathcal{H}_n} \pi(\mathbf{v}_n)$.

Step 4: The type(s) with the highest value of the posterior probability (MAP type) is to be searched out. Since types which do not belong to $\mathcal{H}_n$ have zero posterior probability, a search for the MAP type can be restricted to types which belong to $\mathcal{H}_n$. So, the MAP type $\hat{\mathbf{v}}_n$ is

$$\hat{\mathbf{v}}_n \triangleq \arg \max_{\mathbf{v}_n \in \mathcal{H}_n} P(\text{type} = \mathbf{v}_n | \text{type} \in \mathcal{H}_n)$$

Since, for fixed n and any $\mathbf{v}_n$, $P(\text{type} \in \mathcal{H}_n)$ is a constant, the MAP type turns to be

$$\hat{\mathbf{v}}_n = \arg \max_{\mathbf{v}_n \in \mathcal{H}_n} \pi(\text{type} = \mathbf{v}_n) \quad (1)$$

Thus the MAP type $\hat{\mathbf{v}}_n$ is just the type in $\mathcal{H}_n$ which has the highest value of the prior probability.

Here it stops.

Observe that (1) is identical with prescription of the Maximum Probability (MaxProb) method (cf. [3]). Thus the above reasoning provides its bayesian interpretation.

HOW DOES IT RELATE TO REM/MAXENT?

Via Maximum Probability Theorem (MPT, see [3], Thm 1 and [6]).

Before stating MPT, I-projection has to be defined. I-projection $\hat{\mathbf{p}}$ of $\mathbf{q}$ on set $\Pi \subseteq \mathcal{P}(\mathcal{X})$ is such $\hat{\mathbf{p}} \in \Pi$ that $I(\hat{\mathbf{p}} \| \mathbf{q}) = \inf_{\mathbf{p} \in \Pi} I(\mathbf{p} \| \mathbf{q})$, where¹ $I(\mathbf{p} \| \mathbf{q}) \triangleq \sum_{\mathcal{X}} p_i \log \frac{p_i}{q_i}$ is the I-divergence. I-divergence is known under various other names: Kullback-Leibler's distance, KL number, Kullback's directed divergence, etc. When taken with minus sign it is known as (Shannon-Kullback's) relative entropy.

(MPT)² Let differentiable constraint $F(\mathbf{v}_n) = 0$ define feasible set of types $\mathcal{H}_n$ and let $\mathcal{H} \triangleq \{\mathbf{p} : F(\mathbf{p}) = 0\}$ be the corresponding feasible set of probability mass functions. Let $\hat{\mathbf{v}}_n \triangleq \arg \max_{\mathbf{v}_n \in \mathcal{H}_n} \pi(\mathbf{v}_n)$. Let $\hat{\mathbf{p}}$ be I-projection of $\mathbf{q}$ on $\mathcal{H}$. And let $n \rightarrow \infty$. Then $\hat{\mathbf{v}}_n = \hat{\mathbf{p}}$.

MPT shows that REM is an asymptotic instance of MaxProb method. Thus MPT carries the bayesian interpretation of MaxProb over into REM/MaxEnt. Hence, I-projection is just the MAP type which results from the bayesian updating which was described at the previous Section, in the case of sufficiently large n .

To sum up: Whenever n is sufficiently large and prior will be assigned to types $\mathbf{v}_n$ as in the Step 1, and new data will take form as in the Step 2, and the prior will be updated by the data via BT as in the Step 3, and MAP type will be searched out as in the Step 4, then the MAP type will be nothing but the REM I-projection of $\mathbf{q}$ on $\mathcal{H}$.

DISCUSSION

Why MAP? Why not say median aposteriori type? The MAP type becomes when $n \rightarrow \infty$ just the I-projection. If the I-projection is unique then Conditioned Weak Law of Large Numbers (CWLLN, cf. [13], [12], [7], [1], [11], [9], [10]) can be invoked. If read in the above bayesian manner, it says that any other type/distribution than I-projection has asymptotically zero posterior probability. So, this is

¹ There, $\log 0 = -\infty$, $\log \frac{b}{0} = +\infty$, $0 \cdot (\pm\infty) = 0$, conventions are assumed. Throughout the paper $\log$ denotes the natural logarithm.

² Originally MPT was stated with unique I-projection case in mind. Its proof however readily allows to state it in general form (see [6]). Since the issue of uniqueness is at this Section irrelevant the MPT will be stated at its original form.

why MAP and not median. However, what if there are multiple I-projections? Obviously, the bayesian interpretation of MaxProb is valid regardless of the number of MAP types. MPT in its general form (cf. [6]) covers also the case of multiple MaxProb types and claims that they converge to I-projections. Then one can either recall Entropy Concentration Theorem (cf. [6]) or invoke an extension of CWLLN which covers also the case of multiple I-projections (cf. [5]) – to answer the “Why MAP” question in the general case.

CONCLUDING NOTE

Originally (cf. [3]), MaxProb was presented as a method which looks in $\mathcal{H}_n$ for a type $\hat{\mathbf{v}}_n = \arg \max_{\mathbf{v}_n \in \mathcal{H}_n} \pi(\mathbf{v}_n)$ which the ‘prior’ generator $\mathbf{q}$ can generate with the highest probability. The word ‘prior’ was used merely to mean that the generator is selected before the data arrive. Alternatively, since unconstrained maximization of the conditional probability $P(\text{type} = \mathbf{v}_n | \text{type} \in \mathcal{H}_n)$ reduces to maximization of $\pi(\mathbf{v}_n)$ constrained to $\mathbf{v}_n \in \mathcal{H}_n$, MaxProb could be interpreted as search for the type with the highest value of the conditional probability. The third, bayesian interpretation of MaxProb – inspired by [2] – was given here. Obviously, MPT stands regardless of what is the preferred interpretation of MaxProb.

ACKNOWLEDGEMENTS

Supported by the grant VEGA 1/0264/03 from the Scientific Grant Agency of the Slovak Republic. It is a pleasure to thank Viktor Witkovský for valuable discussions and comments on earlier version of this work. The thanks extend to Aleš Gottvald and George Judge. Lapses are mine.

REFERENCES

1. Csiszár, I., *Ann. Prob.*, **12**, No. 3, 768-793, (1984).
2. Csiszár I., “An extended Maximum Entropy principle and a bayesian justification,” in *Bayesian Statistics 2*, J. M. Bernardo, M. H. DeGroot, D. V. Lindley, A. F. M. Smith (eds.), Elsevier, 83-98, North-Holland, 1985.
3. Grendár, M., Jr. and Grendár, M., “What is the question that MaxEnt answers? A probabilistic interpretation,” in *Bayesian inference and Maximum Entropy methods*, A. Mohammad-Djafari (ed.), 83-94, AIP, Melville (NY), 2001. Also arxiv.org/abs/math-ph/0009020, Sep 2000.
4. Grendár, M., Jr. and Grendár, M., *Entropy*, **3**, 58-63, (2001).
5. Grendár, M., Jr. and Grendár, M., *Acta U. Belii Ser. Math.*, **10**, 3-8, (2003).
6. Grendár, M., Jr. and Grendár, M., “Maximum Entropy method with non-linear moment constraints: challenges,” in *Bayesian inference and Maximum Entropy methods*, G. Erickson and Y. Zhai (eds.), 97-110, AIP, Melville (NY), 2004. Also arxiv.org/abs/physics/0308006.
7. Groeneboom, P., Oosterhoff, J. and Ruymgaart, F. H., *Ann. Prob.*, **7**, 553-586, (1979).
8. Jaynes, E. T., “The relation of bayesian and Maximum Entropy methods,” in *Maximum Entropy and Bayesian Methods in Science and Engineering (Vol. 1)*, G. J. Erickson and C. R. Smith (eds.), Kluwer, pp. 25-29, 1988.
9. La Cour, B. R. and Schieve, W. C., *Jour. Stat. Phys.*, **107**, 3/4, 729-755, (2002).
10. Leonard, Ch. and Najim, J., *Bernoulli*, **8**, 6, 721-743, (2002).
11. Lewis, J. T., Pfister, C.-E. and Sullivan, W. G., *Markov Proc. Rel. Field.*, **1**, 319-386, (1995).
12. van Campenhout, J. M. and Cover, T. M., *IEEE IT*, **27**, 483-489, (1981).
13. Vasicek, O. A., *Ann. Prob.*, **8**, 142-147, (1980).

Corrections wrt Version 1: i) Obviously, $P(\text{type} \in \mathcal{H}_n)$ is not given as a ratio of the number of types in $\mathcal{H}_n$ to the number of all types in Π_n ; rather it is $P(\text{type} \in \mathcal{H}_n) = \sum_{\mathbf{v}^n \in \mathcal{H}_n} \pi(\mathbf{v}^n)$. The rest of argument remains untouched by the gross lapse. ii) The second question from Discussion (Sect. 5) from the Version 1 is not included here.

In the form of Version 1, the paper appeared as M. Grendar, Jr. and M. Grendar, Maximum Probability and Maximum Entropy methods: Bayesian interpretation, in *Bayesian Inference and Maximum Entropy methods in Science and Engineering*, G. Erickson and Y. Zhai (eds.), AIP, Melville, pp. 490 -495, 2004.